

Crop Report


1-Sep-2017

UpperNorthFS: Barrie


Crop: Wheat
Cultivar: Katana
Sowing details: 155 plants/m² on 23-May
Expected maturity date: 21-Nov

Paddock Details
Initial conditions date: 23-May
Soil: Sandy Clay Loam over Light and Medium Clay (Morchard Hill No604-YP)
800 mm max rooting depth
Stubble: 500 kg/ha of Wheat
No till

Grain Yield Outcome




Hay Yield Outcome




Current dry matter: 3727kg/ha

The Season So Far - Growing Season Rainfall Deciles




Simulated and Predicted Crop Growth Stage



Probability and Incidence of Frost and Heat Shock

Frost damage during flowering				Heat damage during grain fill			
Severity	Probability	This Season		Severity	Probability	This Season	
mild 2 to 0°C during flowering	<div></div>	22%	0	mild 32 to 34°C	<div></div>	55%	0
moderate 0 to -2°C during flowering & early grain fill		0%	0	moderate 34 to 36°C	<div></div>	34%	0
severe Less than -2°C during flowering & grain fill		0%	0	severe Above 36°C	<div></div>	22%	0

Current Distribution of PAW




Water Budget

Initial PAW status @ 23-May
Rainfall since 23-May
Irrigations
Evaporation since 23-May
Transpiration since 23-May
Deep drainage since 23-May
Run-off since 23-May

Current PAW status:

Probability of Future Waterlogging Events



Nitrogen Budget

Initial N status @ 23-May
N mineralisation since 23-May
N tie up since 23-May
N applications

24-May : 25 kg/ha
2-Aug : 37 kg/ha


Total N in plant
De-nitrification since 23-May
Leaching since 23-May

Current N status:


Median N mineralisation to maturity = 0.017 kg/ha
Median N tie up to maturity = 1.595 kg/ha

62 kg/ha
1 kg/ha
2 kg/ha
89 kg/ha
0 kg/ha
0 kg/ha
34 kg/ha


Current distribution of soil nitrogen (kg/ha)




Availability of Water to Growing Roots




Availability of Soil Nitrogen to Growing Roots



Water Stress



Nitrogen Stress



Brief periods of mild to moderate stress do not necessarily lead to reduced yield. To see the likely impacts of additional nitrogen fertiliser rates use the Nitrogen and Nitrogen Profit reports.


Median projected crop performance and requirements for the next 10 days assuming no rain and no added fertiliser

Date	Growth Stage	Evap. (mm)	Water use (mm)	N use (kg/ha)	Water avail. to roots above stress threshold (mm)	Water avail. to roots above CLL (mm)	N avail. to roots (kg/ha)	Mineralisation (kg/ha)	N tie up (kg/ha)
1-Sep	33.6	0.6	0.9	0.2	-9.3	13.9	23.7	0.0	0.0
2-Sep	33.9	0.6	0.8	0.2	-10.6	12.6	23.6	0.0	0.0
3-Sep	34.3	0.6	0.7	0.1	-11.9	11.3	23.4	0.0	0.0
4-Sep	34.7	0.6	0.6	0.1	-13.0	10.2	23.2	0.0	0.0
5-Sep	35.1	0.6	0.5	0.1	-14.2	9.0	23.1	0.0	0.0
6-Sep	35.5	0.6	0.5	0.1	-15.3	8.0	23.0	0.0	0.0
7-Sep	36.0	0.5	0.4	0.1	-16.3	6.9	22.9	0.0	0.0
8-Sep	36.3	0.3	0.4	0.1	-17.0	6.2	22.8	0.0	0.0
9-Sep	36.8	0.3	0.3	0.1	-17.7	5.6	22.7	0.0	0.0
10-Sep	37.2	0.3	0.3	0.1	-18.3	5.0	22.7	0.0	0.0


The water available to roots above the stress threshold is the amount of PAW (mm) above one third of the total water holding capacity of this soil. If the water values are below this stress threshold the water available to roots above the stress threshold will be negative.

Bureau of Meteorology Seasonal and Monthly Outlooks


3 MONTH CLIMATE OUTLOOK FROM SEPTEMBER TO NOVEMBER




SEPTEMBER CLIMATE OUTLOOK OCTOBER CLIMATE OUTLOOK




PAST ACCURACY FROM SEPTEMBER TO NOVEMBER



PAST ACCURACY FOR SEPTEMBER



PAST ACCURACY FOR OCTOBER



© Copyright
Commonwealth of Australia 2016 Bureau
of Meteorology
<http://www.bom.gov.au/climate>