



Crop Report


21-Aug-2017

UpperNorthFS: Rodgers

Crop: Wheat
Cultivar: Corack
Sowing details: 110 plants/m² on 2-May
Expected maturity date: 23-Oct


Paddock Details
Initial conditions date: 23-May
Soil: Clay loam over medium clay (Quorn No605)
1800 mm max rooting depth
Stubble: 1000 kg/ha of Wheat
No till

Grain Yield Outcome



This graph shows the probability of exceeding a range of yield outcomes this season. It takes into account your pre-season soil moisture, the weather conditions so far, soil N and agronomic inputs. The long term record from your nominated weather station is then used to simulate what would have happened from this date on in each year of the climate record. The yield results are used to produce this graph.


Hay Yield Outcome




This graph shows the probability of exceeding a range of hay yield outcomes this season. It takes into account the same factors as the grain yield graph above. When above ground dry matter is below 2t/ha, hay yield is assumed to be 70% of dry matter, with a moisture content of 13%. When dry matter is between 2 and 12t/ha, hay yield is assumed to be between 70 and 75% of dry matter (sliding scale). When dry matter is above 12t/ha, hay yield is assumed to be between 75 and 80% (sliding scale).

Current dry matter: 2402kg/ha

The Season So Far - Growing Season Rainfall Deciles




Simulated and Predicted Crop Growth Stage



Probability and Incidence of Frost and Heat Shock

Frost damage during flowering				Heat damage during grain fill			
Severity	Probability	This Season		Severity	Probability	This Season	
mild 2 to 0°C during flowering	<div><div></div></div>	27%	0	mild 32 to 34°C	<div><div></div></div>	29%	0
moderate 0 to -2°C during flowering & early grain fill		0%	0	moderate 34 to 36°C	<div><div></div></div>	18%	0
severe Less than -2°C during flowering & grain fill		0%	0	severe Above 36°C	<div><div></div></div>	3%	0


Current Distribution of PAW



Water Budget

Initial PAW status @ 23-May	69 mm
Rainfall since 23-May	41.1 mm
Irrigations	
Evaporation since 23-May	41 mm
Transpiration since 23-May	38 mm
Deep drainage since 23-May	0 mm
Run-off since 23-May	0 mm
Current PAW status:	36 mm


Probability of Future Waterlogging Events




Nitrogen Budget

Initial N status @ 23-May	38 kg/ha
N mineralisation since 23-May	2 kg/ha
N tie up since 23-May	5 kg/ha
N applications	
25-May :	30.8 kg/ha
Total N in plant	40 kg/ha
De-nitrification since 23-May	0 kg/ha
Leaching since 23-May	0 kg/ha
Current N status:	27 kg/ha
Median N mineralisation to maturity	= 0.741 kg/ha
Median N tie up to maturity	= 3.107 kg/ha


Current distribution of soil nitrogen (kg/ha)




Availability of Water to Growing Roots




Availability of Soil Nitrogen to Growing Roots



Water Stress



Nitrogen Stress



Brief periods of mild to moderate stress do not necessarily lead to reduced yield. To see the likely impacts of additional nitrogen fertiliser rates use the Nitrogen and Nitrogen Profit reports.

Median projected crop performance and requirements for the next 10 days assuming no rain and no added fertiliser

Date	Growth Stage	Evap. (mm)	Water use (mm)	N use (kg/ha)	Water avail. to roots above stress threshold (mm)	Water avail. to roots above CLL (mm)	N avail. to roots (kg/ha)	Mineralisation (kg/ha)	N tie up (kg/ha)
21-Aug	49.1	1.0	1.2	0.5	-5.7	30.6	21.0	0.0	0.1
22-Aug	50.0	0.9	1.0	0.4	-7.6	28.8	20.5	0.0	0.1
23-Aug	50.9	0.4	0.9	0.3	-8.7	27.8	20.2	0.0	0.1
24-Aug	51.7	0.4	1.0	0.3	-9.8	26.7	19.9	0.0	0.1
25-Aug	52.7	0.4	0.9	0.2	-11.0	25.6	19.6	0.0	0.1
26-Aug	53.6	0.4	0.8	0.2	-12.2	24.5	19.4	0.0	0.1
27-Aug	54.4	0.3	0.8	0.2	-13.3	23.4	19.3	0.0	0.1
28-Aug	55.3	0.3	0.8	0.1	-14.2	22.6	19.2	0.0	0.0
29-Aug	56.1	0.3	0.7	0.1	-15.3	21.6	19.1	0.0	0.0
30-Aug	57.0	0.3	0.7	0.1	-16.3	20.6	19.0	0.0	0.0

The water available to roots above the stress threshold is the amount of PAW (mm) above one third of the total water holding capacity of this soil. If the water values are below this stress threshold the water available to roots above the stress threshold will be negative.

Bureau of Meteorology Seasonal and Monthly Outlooks

