

# Crop Report

10-Nov-2017


UpperNorthFS:  
Berryman

Crop: Wheat  
Cultivar: Sceptre  
Sowing details: 150 plants/m<sup>2</sup> on 28-Apr  
Expected maturity date: 3-Nov

Paddock Details  
Initial conditions date: 24-May  
Soil: Red Cracking Clay (Pirie)(CU022)  
1000 mm max rooting depth  
Stubble: 0 kg/ha of Wheat  
No till


## Grain Yield Outcome

- ☒ Nitrogen limited Yield  
☐ Nitrogen limited Yield with Frost and heat Effects  
☒ Water limited Yield  
☐ Water limited Yield with Frost and heat Effects



This graph shows the probability of exceeding a range of yield outcomes this season. It takes into account your pre-season soil moisture, the weather conditions so far, soil N and agronomic inputs. The long term record from your nominated weather station is then used to simulate what would have happened from this date on in each year of the climate record. The yield results are used to produce this graph.


## Hay Yield Outcome




This graph shows the probability of exceeding a range of hay yield outcomes this season. It takes into account the same factors as the grain yield graph above. When above ground dry matter is below 2t/ha, hay yield is assumed to be 70% of dry matter, with a moisture content of 13%. When dry matter is between 2 and 12t/ha, hay yield is assumed to be between 70 and 75% of dry matter (sliding scale). When dry matter is above 12t/ha, hay yield is assumed to be between 75 and 80% (sliding scale).

Current dry matter: 0kg/ha

## The Season So Far - Growing Season Rainfall Deciles




Simulated and Predicted Crop Growth Stage



Probability and Incidence of Frost and Heat Shock

Frost damage during flowering				Heat damage during grain fill			
Severity	Probability	This Season		Severity	Probability	This Season	
mild 2 to 0°C during flowering	<div><div></div></div>	60%	1	mild 32 to 34°C	<div><div></div></div>	27%	0
moderate 0 to -2°C during flowering & early grain fill	<div><div></div></div>	9%	0	moderate 34 to 36°C	<div><div></div></div>	6%	0
severe Less than -2°C during flowering & grain fill	<div><div></div></div>	0%	0	severe Above 36°C	<div><div></div></div>	4%	0


Current Distribution of PAW



Water Budget

Initial PAW status @ 24-May	101 mm
Rainfall since 24-May	97 mm
Irrigations	
Evaporation since 24-May	65 mm
Transpiration since 24-May	140 mm
Deep drainage since 24-May	0 mm
Run-off since 24-May	0 mm
<b>Current PAW status:</b>	<b>3 mm</b>


Probability of Future Waterlogging Events




Nitrogen Budget

Initial N status @ 24-May	161 kg/ha
N mineralisation since 24-May	0 kg/ha
N tie up since 24-May	5 kg/ha
N applications	
25-May : 15 kg/ha	
17-Jul : 35 kg/ha	
1-Aug : 46 kg/ha	
Total N in plant	0 kg/ha
De-nitrification since 24-May	1 kg/ha
Leaching since 24-May	0 kg/ha
<b>Current N status:</b>	<b>54 kg/ha</b>
Median N mineralisation to maturity	- 0 kg/ha
Median N tie up to maturity	- 0 kg/ha


Current distribution of soil nitrogen (kg/ha)




Availability of Water to Growing Roots




Availability of Soil Nitrogen to Growing Roots



Water Stress



Nitrogen Stress




Brief periods of mild to moderate stress do not necessarily lead to reduced yield. To see the likely impacts of additional nitrogen fertiliser rates use the Nitrogen and Nitrogen Profit reports.

Median projected crop performance and requirements for the next 10 days assuming no rain and no added fertiliser


Date	Growth Stage	Evap. (mm)	Water use (mm)	N use (kg/ha)	Water avail. to roots above stress threshold (mm)	Water avail. to roots above CLL (mm)	N avail. to roots (kg/ha)	Mineralisation (kg/ha)	N tie up (kg/ha)
9-Nov	9.0	0.0	0.0	0.0	-15.0	0.0	8.5	0.0	0.0
10-Nov	9.0	0.0	0.0	0.0	-15.0	0.0	8.5	0.0	0.0
11-Nov	9.0	0.0	0.0	0.0	-15.0	0.0	8.5	0.0	0.0
12-Nov	9.0	0.0	0.0	0.0	-15.0	0.0	8.5	0.0	0.0
13-Nov	9.0	0.0	0.0	0.0	-15.0	0.0	8.5	0.0	0.0
14-Nov	9.0	0.0	0.0	0.0	-15.0	0.0	8.5	0.0	0.0
15-Nov	9.0	0.0	0.0	0.0	-15.0	0.0	8.5	0.0	0.0
16-Nov	9.0	0.0	0.0	0.0	-15.0	0.0	8.5	0.0	0.0
17-Nov	9.0	0.0	0.0	0.0	-15.0	0.0	8.5	0.0	0.0
18-Nov	9.0	0.0	0.0	0.0	-15.0	0.0	8.5	0.0	0.0

The water available to roots above the stress threshold is the amount of PAW (mm) above one third of the total water holding capacity of this soil. If the water values are below this stress threshold the water available to roots above the stress threshold will be negative.


3 MONTH CLIMATE OUTLOOK FROM  
OCTOBER TO DECEMBER




OCTOBER  
CLIMATE OUTLOOK



NOVEMBER  
CLIMATE OUTLOOK



PAST ACCURACY FROM  
OCTOBER TO DECEMBER



PAST ACCURACY FOR  
OCTOBER



PAST ACCURACY FOR  
NOVEMBER

